

MINNEAPOLIS ORTHOPAEDICS

Advanced skills and experience for the results you deserve

825 South 8th Street, Suite 550
Minneapolis, MN 55404
Phone – 612.333.5000 Fax – 612.333.6922
Douglas A. Becker, MD
Angela Finn, CNP

HOME CARE AFTER ROTATOR CUFF REPAIR AND LABRAL REPAIR

Wound Care:

- 1) After 3 days remove the dressing.
- 2) Keep a bandage on the wound and change it every 2 days until there is no longer drainage. Some drainage may persist for up to 2 weeks from surgery.
- 3) Apply ice to shoulder for 20 minutes 3 times per day for 2 weeks.
- 4) Keep the wound dry until the stitches are removed.
- 5) Call the office if you develop any of the following:
 - Temperature over 101 degrees
 - Large amount of bleeding from incision
 - Significantly increased redness, pain, numbness, or swelling
 - Thick, yellow, or foul smelling drainage from incision

Sling:

- 1) The sling is to be worn full-time (day and night) for **6 weeks**. You should remove the sling during exercises.

Exercises:

- 1) Perform each directed exercise 3 times per day. Do 10-15 repetitions per exercise. Ice your shoulder after exercises.
- 2) You are not to lift your arm on your own other than directed exercises for 6 weeks. No overhead activity, no lifting, and no moving your arm away from your side.
- 3) Physical Therapy is an important part of your rehabilitation. PT will be arranged when you return for your 2 week post-op appointment if it hasn't already been set-up.

Pain:

- 1) A prescription pain medication will be given to you before you leave the hospital.
- 2) Take 2 aspirin each day for 1 month (unless you are allergic to aspirin or have been directed by another doctor not to take aspirin).
- 3) If you need additional pain medication call the office during office hours.
- 4) Some discomfort in the shoulder is expected for up to 6 weeks. Swelling in the operated shoulder, arm, and hand is often noticeable for up to 4 months.

General Directions:

- 1) Please contact the office, during office hours, to set up your 2 week post-op appointment.
- 2) If you have any questions, concerns, or are unsure of your appointment times call the office during office hours.
- 3) If you are having difficulty sleeping try putting pillow(s) under your surgery shoulder & arm, prop yourself up with pillows, or use a recliner.